

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/272827149>

Environmental controls on C, N and S cycling in plants of the Namib Desert

Conference Paper · April 2011

CITATIONS

0

READS

35

4 authors:



Keir Soderberg

S. S. Papadopoulos & Associates, Inc

32 PUBLICATIONS 691 CITATIONS

SEE PROFILE



Joh R. Henschel

National Research Foundation, South Africa

154 PUBLICATIONS 2,354 CITATIONS

SEE PROFILE



Robert John Swap

NASA

148 PUBLICATIONS 3,668 CITATIONS

SEE PROFILE



Stephen A. Macko

University of Virginia

401 PUBLICATIONS 27,532 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Infrared Based Isotope Ratio Measurements for 15N [View project](#)



Ecohydrology of roots [View project](#)



Environmental controls on C, N and S cycling in plants of the Namib Desert

Keir Soderberg (1,2), Joh Henschel (3), Robert Swap (2), and Stephen Macko (2)

(1) Department of Civil and Environmental Engineering, Princeton University, Princeton, NJ, USA (soderbrg@princeton.edu),
(2) Department of Environmental Sciences, University of Virginia, Charlottesville, VA, USA, (3) Gobabeb Training and Research Centre, Walvis Bay, Namibia

Nutrient cycles (C, N and S) in the Namib Desert are investigated through a survey of $\delta^{15}\text{N}$, $\delta^{13}\text{C}$ and $\delta^{34}\text{S}$ of plant material and soils along a fog gradient. The coastal Namib Desert in southwestern Africa is hyper-arid in terms of rainfall, but receives up to 100 days of fog each year. This climate regime leads to interesting water relations among the desert flora and fauna. Among many enigmatic characteristics, photosynthesis in *W. mirabilis* has puzzled researchers since the 1970's. Although it is predominantly a C3 plant, $\delta^{13}\text{C}$ ranges from -17.4 to -24.2 ‰ in natural habitats, and can be as enriched as -14.4 ‰ under artificial growing conditions. Recently the CAM pathway has been confirmed, but it is dominated by CAM-cycling rather than primary CAM fixation, and the driver for CAM utilization has not been identified. Literature values of $\delta^{13}\text{C}$ are replotted across a 150 km aridity gradient for comparison with new data from Hope Mine, showing a significant correlation that appears to be explained by rainfall ($R^2 = 0.25$, $p < 0.001$). Within the Hope Mine community, there are significant positive relationships among stem size, N, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$. The responsible mechanism needs further investigation, but high levels of CAM-cycling and photorespiration along with recycling of N could explain the observed patterns. In addition, there appears to be a reversal of the trend between rainfall and $\delta^{15}\text{N}$ found in Kalahari C3 plants, as more depleted $\delta^{15}\text{N}$ values were found closer to the coast.