

Stock Structure of Snoek *Thyrsites Atun* in the Benguela: A New Hypothesis

M. H. Griffiths

To cite this article: M. H. Griffiths (2003) Stock Structure of Snoek *Thyrsites Atun* in the Benguela: A New Hypothesis, African Journal of Marine Science, 25:1, 383-386, DOI: [10.2989/18142320309504025](https://doi.org/10.2989/18142320309504025)

To link to this article: <https://doi.org/10.2989/18142320309504025>



Published online: 08 Jan 2010.



Submit your article to this journal [↗](#)



Article views: 270



Citing articles: 5 View citing articles [↗](#)

STOCK STRUCTURE OF SNOEK *THYRSITES ATUN* IN THE BENGUELA: A NEW HYPOTHESIS

M. H. GRIFFITHS*

It has long been accepted that snoek *Thyrsites atun* in the Benguela system constitute a single stock that undergoes seasonal longshore migration in waters between southern Angola and the west coast of South Africa. Based on a review of past literature and evaluation of new data, it is contended that Benguela snoek exist rather as two separate sub-populations – divided by the upwelling cell in southern Namibia – with limited, medium-term exchange (c. 5 years), driven mainly by temperature and food availability.

Key words: migration, nursery areas, spawning grounds, sub-populations, *Thyrsites atun*

The snoek *Thyrsites atun* is a medium-sized, pelagic predator (max. size 9 kg, Nepgen 1979a) that inhabits the coastal waters of the temperate southern hemisphere (Kailola *et al.* 1993). Southern African snoek have been recorded from northern Angola to Algoa Bay on the South African east coast, but are found mostly between the Kunene River and Cape Agulhas, i.e. in the Benguela ecosystem. *T. atun* has been an important commercial species in the system since the early 1800s, caught initially with handlines but also by trawlers after 1960 (Crawford 1995). The total reported catch peaked at about 81 000 tons in 1978, but it then dropped considerably following the exclusion of foreign trawlers from Namibian fishing grounds in 1991 (FAO 1978, 1981, 1990, 1997). Between 1991 and 1994 the annual catch ranged between 14 437 and 22 920 tons, of which 93% was taken in South African waters (FAO 1995). *T. atun* is by far the most important species caught by the South African commercial linefishery (constituting 39% of catches between 1986 and 1997). It is also targeted by recreational anglers, but catch statistics are not available for this fishing sector. Around 40% of the reported South African catch between 1990 and 1996 was made by commercial handline fishers and 60% by trawlers. *T. atun* is one of the major predators of anchovy *Engraulis capensis* and sardine *Sardinops sagax* in the southern Benguela (Wickens *et al.* 1992), and it has been implicated in top-down effects on both prey and consequently zooplankton populations (Verheye *et al.* 1998).

Despite its circumglobal distribution, *T. atun* is confined to coastal waters where it often consists of discrete stocks. For example, three stocks are recognized off New Zealand (Hurst and Bagley 1989) and there are between three and five stocks off Australia (Blackburn and Gartner 1954, Grant *et al.* 1978). Understanding

the stock structure of snoek in the Benguela system is essential to successful and realistic ecosystem modelling as well as wise management of the resource.

Prompted by large interannual fluctuations in *T. atun* availability, and the profound effects these had on the linefishery, the first investigation into their migratory patterns and stock structure in the Benguela was conducted in 1934, involving the tagging of 3 755 fish off Namibia (22–23°S). Of these 17 were recaptured, and all had moved southwards, 13 (74%) into South African waters (De Jager 1955). Some of these 13 fish moved as far south as Cape Point off the Cape Peninsula, a distance of about 1 300 km. Based on these results, on anecdotal reports on the seasonal nature of line catches at different localities (Van Wyk 1944, Davies 1954, De Jager 1955), and on temporal patterns in trawl catches, Crawford and De Villiers (1985) postulated that snoek in the Benguela constituted a single stock. The population was believed to undergo a seasonal longshore migration, moving south into South African waters to spawn in winter, then returning north, to as far as southern Angola, in spring/summer. This theory became widely accepted (Crawford *et al.* 1987, Crawford 1995), but a recent study by Griffiths (2002) showed that adult snoek are available to South African linefishers throughout the year, and that the seasonal availability of adults on the trawl grounds results from an offshore spawning migration rather than southward movement from Namibian waters. Griffiths (2002) also demonstrated that higher handline catch rates north of Cape Columbine (Fig. 1) in autumn were caused by juvenile snoek following clupeoid recruits inshore (i.e. within range of linefish vessels) rather than the southward displacement of adults en route to the Agulhas Bank. The results of a South African tagging study conducted during the mid 1970s (3 139 tagged

* Marine & Coastal Management, Private Bag X2, Rogge Bay 8012, Cape Town, South Africa. E-mail: mgriffit@mcm.wcape.gov.za

Manuscript received March 2002; accepted May 2002




Fig. 1: Conceptual model summarizing available knowledge of the life history of Benguela snoek and prevailing current patterns. South African life history information after Griffiths (2002); Namibian life history information inferred from spatial patterns in size composition presented by Vendiktova (1987) egg and larval distributions after Olivar and Fortuño and Olivar and Shelton; and physical oceanography after Boyd *et al.* (1992) and Gründlingh (1999)

and 18 recaptured) are also not consistent with the concept of seasonal longshore movement of adult snoek; despite liberty periods of 22–377 days (seven for >200 days), only local and random movement was recorded (Nepgen 1979a).

Ichthyoplankton surveys conducted between 1977 and 1986 reveal that snoek eggs and larvae are present throughout the Benguela system in winter/spring, distributed as two disjunct bands (Fig. 1) that are separated by the intense Lüderitz upwelling cell (25–27°S) off southern Namibia (Olivar and Fortuño 1991, Olivar and Shelton 1993). The results indicate that spawning

occurs simultaneously in both northern and southern Benguela, which negates the spawning migration hypothesis, and furthermore implies that the southern African snoek stock may exist as two separate sub-populations. The Lüderitz upwelling cell represents a major environmental barrier to many fish species, including anchovy and sardine (Boyd and Hewitson 1983, Cruickshank and Boyd 1985), principal prey items of *T. atun* (Nepgen 1979b, 1982, Griffiths 2002). The concept of two separate sub-populations is further supported by:

- (i) low snoek catches in the vicinity of the environmental barrier (Crawford and De Villiers 1985);
- (ii) the existence of spawning grounds and nursery areas in both South African (Griffiths 2002) and Namibian (Vendiktova 1987) waters (Fig. 1); and
- (iii) the local and random movement of adult snoek tagged off South Africa.

An inverse relationship between annual handline catch trends off Namibia and South Africa (Crawford *et al.* 1990, 1995) suggests migration between sub-populations in the medium term (approximately 5 years), but that not all fish migrate from the waters of one country to those of the other. Given that snoek biomass off each of these countries has been correlated with the abundance of principal prey species (Crawford 1995), it may be that limited medium-term migration occurs in response to prey availability.

A model of separate sub-populations, with medium-term migration, also accommodates the “contradictory” results of the two tagging studies. The tagging experiment conducted by Nepgen (1979a) between 1973 and 1975 took place when handline catches off South Africa were high and off Namibia were low (Crawford *et al.* 1995). This was apparently the result of reduced prey availability in Namibia, caused by the southward intrusion of warm equatorial water, following a Benguela *Niño* (Shannon *et al.* 1988), and a coincident increase of anchovy off South Africa (Shelton *et al.* 1985). Therefore, it is not surprising that no migration to the northern sub-population was recorded. On the other hand, the 1934 tagging experiment off Namibia was conducted during a major Benguela *El Niño* event (Shannon *et al.* 1986), which resulted in the southward movement of all recaptured fish. Although it may be contended that migratory patterns could switch from extensive seasonal movement to stock displacement (Crawford *et al.* 1990), the facts that large fluctuations in South African handline catch have taken place in accordance with prey availability since the late 1800s (Gilchrist 1914, Crawford *et al.* 1987), and that catches off South Africa and Namibia have been negatively correlated since as early as 1965 (the earliest date for which catch data are available for both countries), to-

gether tend to render this argument unlikely.

It is possible that three genetically distinct populations exist: two resident and one migratory (*sensu* Fréon and Misund 1999). Even though catch statistics reveal that not all snoek migrate, genetic (satellite DNA) and/or extensive tagging studies would be necessary to test this hypothesis. However, regardless of whether there are two sub-populations or three distinct stocks, interannual (v. seasonal) movement patterns are not common in pelagic fish. Moreover, examples of "interannual variation of habitat selection" (Fréon and Misund 1999) involve changes to migratory routes rather than the same route with a multi-year cycle.

The stock structure and movement patterns of *T. atun* in other parts of the southern hemisphere are complex. For example, Australian populations appear to overlap spatially but not temporally (Blackburn and Gartner 1954). In addition, some stocks undertake extensive seasonal migrations of several hundred kilometres, e.g. Australia and the east coast of New Zealand, whereas others undergo only local movement, e.g. Chatham Islands (New Zealand) and Tristan da Cunha (Blackburn and Gartner 1954, Hurst and Bagley 1989, Andrew *et al.* 1995). The existence of sub-populations off southern Africa, which mix in the medium term through extensive migration, provides additional variation to an already diverse theme.

Although snoek found off Namibia and South Africa appear to be separate sub-populations, the fact that there may be extensive interaction in the medium term means that Benguela snoek are ultimately a shared resource. The implications of the sub-population model for international management are, however, extremely complex. For example, the impact of fishing activity off one country on the catches of the other will vary according to the temporal proximity of the catch date to a migratory event. Given that movement between sub-populations may be driven by fluctuations in prey availability, and that biomass predictions for small pelagic fish (i.e. snoek prey) are unreliable, long-range predictions of snoek migration are not possible at this stage. Assuming that dynamic properties would vary between sub-populations (in accordance with environmental conditions), future stock assessments should, while treating the resource as a single stock, incorporate information from both sub-populations. Attempts should also be made to standardize management objectives, biological reference points and regulations (e.g. minimum size) through a bi-national management plan.

Although southward movement of snoek from Namibian to South African waters has been confirmed by tag recaptures, northward movement between the two countries has not been established. It is recom-

mended that tagging programmes be implemented simultaneously in both countries to provide a better understanding of rates of exchange between sub-populations.

LITERATURE CITED

- ANDREW, T. G., HECHT, T., HEEMSTRA, P. C. and J. R. E. LUTJEHARMS 1995 — Fishes of the Tristan da Cunha Group and Gough Island, South Atlantic. *Ichthyol. Bull. J. L. B. Smith Inst. Ichthyol.* **63**: 43 pp.
- BLACKBURN, M. and P. E. GARTNER 1954 — Populations of barracouta *Thyrsites atun* (Euphrasen) in Australian waters. *Aust. J. mar. Freshwat. Res.* **5**: 411–468.
- BOYD, A. J. and J. D. HEWITSON 1983 — Distribution of anchovy larvae off the west coast of southern Africa between 32°30' and 26°30' S, 1979–1982. *S. Afr. J. mar. Sci.* **1**: 71–75.
- BOYD, A. J., TAUNTON-CLARK, J. and G. P. J. OBERHOLSTER 1992 — Spatial features of the near-surface and midwater circulation patterns off western and southern South Africa and their role in the life histories of various commercially fished species. In *Benguela Trophic Functioning*. Payne, A. I. L., Brink, K. H., Mann, K. H. and R. Hilborn (Eds). *S. Afr. J. mar. Sci.* **12**: 189–206.
- CRAWFORD, R. J. M. 1995 — Snoek and chub mackerel. In *Oceans of Life off Southern Africa*, 2nd ed. Payne, A. I. L. and R. J. M. Crawford (Eds). Cape Town: Vlaeberg: 177–187.
- CRAWFORD, R. J. M. and G. DE VILLIERS 1985 — Snoek and their prey – interrelationships in the Benguela upwelling system. *S. Afr. J. Sci.* **81**(2): 91–97.
- CRAWFORD, R. J. M., SHANNON, L. J. and G. NELSON 1995 — Environmental change, regimes and middle-sized pelagic fish in the South-East Atlantic Ocean. In *International Symposium on Middle-Sized Pelagic Fish*. Bas, C., Castro, J. J. and J. M. Lorenzo (Eds). *Scientia Mar., Barcelona* **59**: 417–426.
- CRAWFORD, R. J. M., SHANNON, L. V. and D. E. POLLOCK 1987 — The Benguela ecosystem. 4. The major fish and invertebrate resources. In *Oceanography and Marine Biology. An Annual Review* **25**. Barnes, M. (Ed.). Aberdeen: University Press: 353–505.
- CRAWFORD, R. J. M., UNDERHILL, L. G. and J. D. VENTER 1990 — Handline catches and stock identity of snoek *Thyrsites atun* off South Africa and Namibia. *S. Afr. J. mar. Sci.* **9**: 95–99.
- CRUICKSHANK, R. A. and A. J. BOYD 1985 — An environmental basin model for West Coast pelagic fish distribution. In *International Symposium on the Most Important Upwelling Areas off Western Africa (Cape Blanco and Benguela)*, [Barcelona, 1983] I. Bas, C., Margalef, R. and P. Rubiés (Eds). Barcelona; Instituto de Investigaciones Pesqueras: 541–549.
- DAVIES, D. H. 1954 — The South African pilchard (*Sardinops ocellata*). Development, occurrence and distribution of eggs and larvae, 1950–51. *Invest. Rep. Div. Fish. S. Afr.* **15**: 28 pp.
- DE JAGER, B. VAN D. 1955 — The South African pilchard (*Sardinops ocellata*). The development of the snoek (*Thyrsites atun*) a fish predator of the pilchard. *Invest. Rep. Div. Fish. S. Afr.* **19**: 16 pp.
- FAO 1978 — Yearbook of fishery statistics. Catch and landings, 1977. *FAO Fish Ser.* **46**: 128 pp.
- FAO 1981 — Yearbook of fishery statistics. Catches and landings, 1980. *FAO Fish. Ser.* **52**: 386 pp.
- FAO 1990 — Yearbook of fishery statistics. Catches and landings, 1989. *FAO Fish. Ser.* **70**: 325 pp.

- FAO 1995 — Yearbook of fishery statistics. Catches and landings, 1994. *FAO Fish. Ser.* **80**: 351 pp.
- FAO 1997 — Yearbook of fishery statistics. Catches and landings, 1996. *FAO Fish. Ser.* **84**: 343 pp.
- FRÉON, P. and O. A. MISUND 1999 — *Dynamics of Pelagic Fish Distribution and Behaviour: Effects on Fisheries and Stock Assessment*. London; Fishing News Books: 348 pp.
- GILCHRIST, J. D. F. 1914 — The snoek and allied fishes in South Africa. *Mar. Biol. Rep., Cape Tn* **2**: 116–127.
- GRANT, C. J., COWPER, T. R. and D. D. REID 1978 — Age and growth of snoek, *Leionura atun* (Euphrasen), in south-eastern Australian waters. *Aust. J. mar. Freshwat. Res.* **29**: 435–444.
- GRIFFITHS, M. H. 2002 — Life history of South African snoek *Thyrsites atun* (Pisces: Gempylidae): a pelagic predator of the Benguela ecosystem. *Fishery Bull., Wash.* **100**(4).
- GRÜNDLINGH, M. L. 1999 — Surface currents derived from satellite-tracked buoys off Namibia. *Deep-Sea Res. II* **46**: 453–473.
- HURST, R. J., and N. W. BAGLEY 1989 — Movements and possible stock relationships of the New Zealand barracouta, *Thyrsites atun*, from tag returns. *N. Z. J. mar. Freshwat. Res.* **23**: 105–111.
- KAILOLA, P. J., WILLIAMS, M. J., STEWART, P. C., REICHEL, R. E., McNEE, A. and C. GRIEVE 1993 — *Australian Fisheries Resources*. Canberra; Bureau of Resource Sciences and the Fisheries Research and Development Corporation: 422 pp.
- NEPGEN, C. S. DE V. 1979a — The food of the snoek *Thyrsites atun*. *Fish. Bull. S. Afr.* **11**: 39–42.
- NEPGEN, C. S. DE V. 1979b — Trends in the line fishery for snoek *Thyrsites atun* off the South-Western Cape, and in size composition, length-weight relationship and condition. *Fish. Bull. S. Afr.* **12**: 35–43.
- NEPGEN, C. S. DE V. 1982 — Diet of predatory and reef fish in False Bay and possible effects of pelagic purse-seining on their food supply. *Fish. Bull. S. Afr.* **16**: 75–93.
- OLIVAR, M-P. and J-M. FORTUÑO 1991 — Guide to ichthyoplankton of the Southeast Atlantic (Benguela Current region). *Scientia Mar., Barcelona* **55**(1): 383 pp.
- OLIVAR, M-P. and P. A. SHELTON 1993 — Larval fish assemblages of the Benguela Current. In *Advances in the Early Life History of Fish. 1. Larval Fish Assemblages and Ocean Boundaries*. Moser, H. G., Smith, P. E. and L. A. Fuiman (Eds). *Bull. mar. Sci.* **53**(2): 450–474.
- SHANNON, L. V., BOYD, A. J., BRUNDRIT, G. B. and J. TAUNTON-CLARK 1986 — On the existence of an *El Niño*-type phenomenon in the Benguela system. *J. mar. Res.* **44**(3): 495–520.
- SHANNON, L. V., CRAWFORD, R. J. M., BRUNDRIT, G. B. and L. G. UNDERHILL 1988 — Responses of fish populations in the Benguela ecosystem to environmental change. *J. Cons. perm. int. Explor. Mer* **45**(1): 5–12.
- SHANNON, L. V. and G. NELSON 1996 — The Benguela: large scale features and processes and system variability. In *The South Atlantic: Present and Past Circulation*. Wefer, G., Berger, W. H., Siedler, G. and D. J. Webb (Eds). Berlin; Springer: 163–210.
- SHELTON, P. A., BOYD, A. J. and M. J. ARMSTRONG 1985 — The influence of large-scale environmental processes on neritic fish populations in the Benguela Current system. *Rep. Calif. coop. oceanic Fish. Invest.* **26**: 72–92.
- VAN WYK, G. F. 1944 — South African fish products. 8. Composition of the flesh of Cape fishes. *J. Soc. chem. Ind. Lond.* **63**: 367–371.
- VENIDIKTOVA, L. I. 1987 — Peculiarities of snoek (*Thyrsites atun* Euphr., 1791) distribution in the Namibian area. *Colln scient. Pap. int. Commn SE. Atl. Fish.* **14**(2): 283–286.
- VERHEYE, H. M., RICHARDSON, A. J., HUTCHINGS, L., MARSKA, G. and D. GIANOKOURAS 1998 — Long-term trends in the abundance and community structure of coastal zooplankton in the southern Benguela system, 1951–1996. In *Benguela Dynamics: Impacts of Variability on Shelf-Sea Environments and their Living Resources*. Pillar, S. C., Moloney, C. L., Payne, A. I. L. and F. A. Shillington. *S. Afr. J. mar. Sci.* **19**: 317–332.
- WICKENS, P. A., JAPP, D. W., SHELTON, P. A., KRIEL, F., GOOSEN, P. C., ROSE, B., AUGUSTYN, C. J., BROSS, C. A. R., PENNEY, A. J. and R. G. KROHN 1992 — Seals and fisheries in South Africa – competition and conflict. In *Benguela Trophic Functioning*. Payne, A. I. L., Brink, K. H., Mann, K. H. and R. Hilborn (Eds). *S. Afr. J. mar. Sci.* **12**: 773–789.